

Provided for non-commercial research and education use.
Not for reproduction, distribution or commercial use.



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/authorsrights>



Facile, efficient and eco-friendly synthesis of 5-sulfenyl tetrazole derivatives of indoles and pyrroles

Bruna L. Kuhn ^a, Margiani P. Fortes ^a, Teodoro S. Kaufman ^{b,*}, Claudio C. Silveira ^{a,*}

^a Departamento de Química, Universidade Federal de Santa Maria, 97105-900 Santa Maria, RS, Brazil

^b Instituto de Química Rosario (IQUIR, CONICET-UNR) and FCBYF—Universidad Nacional de Rosario, Suipacha 531, 2000 Rosario, Argentina

ARTICLE INFO

Article history:

Received 24 December 2013

Revised 20 January 2014

Accepted 22 January 2014

Available online 31 January 2014

ABSTRACT

A concise, two-step eco-friendly approach towards the synthesis of 5-sulfenyl tetrazole derivatives of indoles and pyrroles, is reported. The synthesis comprises the oxone-mediated thiocyanation of the starting heterocycles towards intermediate 3-thiocyanato indoles and 2-thiocyanato pyrroles, and their subsequent treatment with sodium azide in 2-propanol/water under zinc bromide promotion.

© 2014 Elsevier Ltd. All rights reserved.

Keywords:

5-Sulfenyl tetrazoles

Pyrrole and indole derivatives

Thiocyanation

Eco-friendly functionalization of heterocycles

Zinc bromide promoted reaction

The tetrazoles are five-membered ring 6 π -heterocycles, containing four contiguous nitrogen atoms.^{1a} They are not found in Nature;^{1b} however, tetrazole derivatives possess interesting applications in agriculture, pharmacy, photography and as components of explosives. In medicinal chemistry, they are employed as metabolically-stable carboxylic acid isosteres.²

5-Substituted tetrazoles are usually prepared by a [3+2] cycloaddition reaction between organic nitriles and hydrazoic acid,³ commonly generated *in situ* from azides, including NaN₃,⁴ TMSN₃,^{5a,b} tin^{5c,d} and organoaluminium azides,^{5e,f} and DPPA,^{5g} and protic or Lewis acids such as I₂,^{6a} NaHSO₄/SiO₂,^{6a} TMSCl,^{6b} Et₃N-HCl,^{6c} NH₄Cl,^{7a} NH₄Cl/LiCl,^{7b} Cu₂O,^{7c} AlCl₃^{7d} and FeCl₃.^{4a} An intramolecular version has also been reported, employing an organic azide moiety.⁸

More recently, the use zinc promoters, especially ZnBr₂ in water-based solvents, has greatly simplified the synthesis.⁹ Variations including mesoporous ZnS nanospheres,^{10a} Zn/Al hydroxalite,^{10b} Bu₂SnO^{10c} and ZnCl₂ under solventless conditions have also been reported.^{10d} Ionic liquids^{10e} and a catalyst-free method in glycerol^{10f} have been proposed as alternative media.

Thiocyanates are versatile chemical intermediates, which embody a strategically useful entry to various derivatives carrying C-S bonds.¹¹ Thiocyanation of heteroaromatics has been carried

out by reaction with inorganic thiocyanate salts under assistance by different acid catalysts or oxidants. Acids include montmorillonite K-10,^{12a} acid alumina,^{12b} TsOH^{12c} and Amberlyst-15,^{12d} while oxidants encompass HIO₃,^{13a} I₂O₅,^{13b} I₂/MeOH,^{13c} CAN,^{13d} Mn(OAc)₃,^{13e} FeCl₃,^{13f} DDQ,^{13g} H₂O₂^{13h} and Oxone[®],¹³ⁱ among others.

5-Sulfenyl tetrazoles are a small and special group of tetrazoles that can be found as a distinguished motif in active pharmaceutical ingredients, such as cefamandol and latamoxef, and other potentially bioactive molecules (Fig. 1).¹⁴

5-Sulfenyl tetrazoles have been occasionally synthesized from thiocyanates and inorganic azides during studies on 5-substituted tetrazoles.^{7a,b,9a}

In addition, 1-substituted 5-thiotetrazoles have been used as capping agents for stabilization of gold nanoparticles,^{15a} while 5-ethyl- and 5-benzyl-thio-1H-tetrazole are used as 1H-tetrazole replacements in oligonucleotide synthesis.^{15b,c} The sulfur atom of the thioether group conveys better solubility and turns the tetrazole ring more acidic than the corresponding 5-alkyl tetrazoles, improving their ability to act as activators.¹⁶

NH₄Cl^{15,17} and ZnBr₂^{4b} have been employed as promoters for the synthesis of 5-thio- and 5-seleno tetrazoles; Et₃N-HCl^{18a} and magnetic Fe₃O₄ nanoparticles have also been used recently.^{18b} Additionally, 5-alkyl(aryl)thiotetrazoles, were advantageously prepared in a toluene-water biphasic system under phase-transfer catalysis which eased work-up operations,^{17b} and further, Et₂AlN₃ was also used as a source of the azide ion.^{5e,f}

* Corresponding authors. Tel./fax: +55 55 3220 8754.

E-mail addresses: kaufman@iquir-conicet.gov.ar (T.S. Kaufman), silveira@quimica.ufsm.br (C.C. Silveira).





Figure 1. Examples of bioactive 5-sulfenyl tetrazoles.



Scheme 1. Proposed reaction sequence towards 5-sulfenyl tetrazoles.

Taking into account our interest in the development of new approaches to the synthesis of polysubstituted nitrogen heterocycles with potential as bioactive compounds, herein we wish to report the synthesis of 5-sulfenyl tetrazole derivatives of indoles and pyrroles in a two-step sequence involving the thiocyanation of the substituted heterocycles and the subsequent reaction of the resulting thiocyanates with NaN_3 under ZnBr_2 promotion, as shown in **Scheme 1**.

The different starting indoles (**1**) were synthesized according to the literature.¹⁹ The intermediate thiocyanates **3** and **4** were prepared by reaction of the indoles and *N*-phenyl pyrroles (**2**)²⁰ with

NH_4SCN , employing oxone as oxidizing agent. Following an adaptation of the work of Wu¹³ⁱ and based on our previous results,²⁰ the amount of NH_4SCN was limited to 1 mmol/mmol of heterocycle in order to avoid formation of polythiocyanated compounds.

In all cases (**Table 1**), the thiocyanation was efficient and furnished the expected compounds in short times and with high regioselectivity. Only 3-substituted indoles and 2-substituted pyrroles were obtained.²¹ In their infrared spectra, all compounds exhibited the typical CN stretching band of the SCN moiety at 2100–2200 cm^{-1} and a resonance at 110–113 ppm in the ^{13}C NMR spectra. The mechanism of the thiocyanation has been studied.¹³ⁱ

With the thiocyanato derivatives in hand, the next step was the [3+2] cycloaddition towards the tetrazoles. This was attempted under different conditions and later optimized, by systematically changing the solvent, amount of NaN_3 and promoter. The effects of time and reaction temperature were also studied, with the results summarized in **Table 2**.


Employing the protocols of the groups of Das^{6a} (entry 1) and Hallberg,²² low product yields and incomplete consumption of the starting material were observed (entry 2). On the other hand, use of glycerol as the reaction medium without promoter (entry 3)^{10f} and water as the sole solvent^{9a} gave no product (entry 4). However, the approach reported by the group of Suresh babu^{4b} met with better luck, furnishing 89% of the expected heterocycle (entry 5). Apparently, in this case the addition of 2-propanol was a key for the success of transformation.

Based on this result, it was decided to investigate the substitution of ZnBr_2 with ZnCl_2 , observing that this caused an increase of the reaction time to 7 h, while reducing the yields to 53% (entry 6). A decrease in the amount of solvent (from 30 to 10 mL) allowed decreasing the reaction time while maintaining the yields (entry 7). Lowering the amount of NaN_3 to 1.2 equiv and further decreasing the amount of the solvent mixture to 6 mL/mmole of thiocyanate (entry 8) resulted in 94% yield of the tetrazole, after 2 h. However, reducing the amount of promoter and running the reaction at 70 °C or room temperature (entries 9–11) proved non-satisfactory.

These tests demonstrated the need of stoichiometric amounts of the promoter and high temperatures for satisfactory results

Table 1

Synthesis of thiocyanates **3** and **4** by reaction of indoles **1** and pyrroles **2** with NH_4SCN and Oxone® in MeOH




Entry No	R	R_1	R_2	R_3	Prod. No	Time (h)	Yield ^a (%)
1	H	H	H	—	3a	1	93
2	5-Br	H	H	—	3b	3	90
3	5-(<i>p</i> -Tolyl)	H	H	—	3c	4	80
4	H	H	Me	—	3d	2	91
5	H	Me	H	—	3e	2	96
6	H	<i>n</i> -C ₈ H ₁₇	H	—	3f	5	95
7	H	Ph	H	—	3g	3.5	90
8	Br	Me	H	—	3h	3	95
9	—	—	—	H	4a	2	89
10	—	—	—	4-OMe	4b	0.5	92
11	—	—	—	4-Cl	4c	1.5	92
12	—	—	—	3-Cl	4d	1.5	90
13	—	—	—	2-Cl	4e	1.5	90

^a Isolated yields after column chromatography.

Table 2

Optimization of the synthesis of the 5-sulfonyl tetrazoles



Entry No	Solvent (mL)	NaN ₃ (equiv)	Promoter (equiv)	Temp (°C)	Time (h)	Yield ^a (%)
1	DMF (1.5)	1.5	I ₂ (0.06)	120	4	76
2	DMF (1.0)	1.1	NH ₄ Cl (1.2)	160	0.25	64 ^b
3	Glycerol (20)	1.2	—	110	6	—
4	H ₂ O (10)	1.1	ZnBr ₂ (1.0)	Reflux	5	—
5	H ₂ O/2-PrOH (1:1, 30)	2.5	ZnBr ₂ (1.0)	Reflux	5	89
6	H ₂ O/2-PrOH (1:1, 10)	2.5	ZnCl ₂ (1.0)	Reflux	7	53
7	H ₂ O/2-Pr ₂ (1:1, 10)	2.5	ZnBr ₂ (1.0)	Reflux	4	87
8	H ₂ O/2-PrOH (1:1, 6)	1.2	ZnBr ₂ (1.0)	Reflux	2	94
9	H ₂ O/2-PrOH (1:1, 6)	1.2	ZnBr ₂ (0.5)	Reflux	3	78
10	H ₂ O/2-PrOH (1:1, 6)	1.2	ZnBr ₂ (1.0)	r.t.	3	39
11	H ₂ O/2-PrOH (1:1, 6)	1.2	ZnBr ₂ (1.0)	70	3	64

^a Isolated yields after column chromatography.^b Performed under microwave irradiation.

and defined the optimum conditions as those employed in entry 8.²³

In order to assess the scope and limitations of the transformation, the 3-thiocyanato-1*H*-indoles prepared as shown in Table 1 were submitted to the optimized conditions, with the results displayed in Table 3. All tetrazole products are novel. Their structures were fully characterized by ¹H and ¹³C NMR analysis, FT-IR spectroscopy and elemental analysis or HRMS.²⁴ The tetrazole carbon atoms displayed a characteristic signal in a very narrow region (δ 154–156 ppm) of the ¹³C NMR spectrum.


It was observed that the less substituted heterocycle (entry 1) reacted more rapidly than its congeners. On the other hand, substituents exert a large influence. For the *N*-substituted compounds, **3e** reacted faster than **3f** but furnished comparatively lower yields of product (entries 5 and 6). However, introduction of a 5-Br substituent improved the yields at the expense of an increase in the reaction time (entries 5 and 8), compared with the non-methylated analogue **3c** (entry 2).

For the series of tetrazoles derived from 1-aryl-2-thiocyanato-1*H*-pyrroles (**4a–e**), the reaction times were generally shorter

and the yields remained good to excellent. No significant steric (entry 13) nor electronic (entries 9–11) effects were observed.

The reaction mechanism for the synthesis of tetrazoles has been discussed elsewhere. Pioneering research by Himo et al.,^{25a} based on computational and experimental studies, demonstrated the role of Zn in the synthesis of tetrazoles. Using MeCN, MeN₃ and ZnBr₂, they were able to prove that there is a significant decrease in the activation energy of the reaction when the nitrogen of the nitrile group is coordinated to the Lewis acid. Whereas binding of Zn^{II} to the azide ion gave calculated activation energy of 34 and 36 kcal mol⁻¹ for tetrahedral and octahedral coordinating zinc ions, respectively, when Zn^{II} was connected to the nitrogen atom of the nitrile, the activation energy dropped to 25–30 kcal mol⁻¹. Activation of the nitrile substrate is also the basis of the mode of action of a new organocatalyst for the synthesis of tetrazoles.^{6b}

In conclusion, a two step synthesis of new 5-thio-tetrazoles derived from 3-thiocyanato-1*H*-indoles and 2-thiocyanato-1-phenyl pyrroles, is reported. The NH₄SCN/oxone reagent combination was employed for the thiocyanation stage, affording good to excellent yields of the thiocyanate intermediates. The [3+2]

Table 3Synthesis of 5-sulfonyl tetrazoles **5** and **6** by reaction of 3-indolyl (**3**) and 2-pyrrolyl (**4**) thiocyanates

Entry No	R	R ₁	R ₂	R ₃	St. mat./prod. no	Time (h)	Yield ^a (%)
1	H	H	H	—	3a/5a	2	94
2	5-Br	H	H	—	3b/5b	3.5	92
3	5-(<i>p</i> -Tolyl)	H	H	—	3c/5c	4	87
4	H	H	Me	—	3d/5d	4	87
5	H	Me	H	—	3e/5e	3	83
6	H	<i>n</i> -C ₈ H ₁₇	H	—	3f/5f	4.5	95
7	H	Ph	H	—	3g/5g	4	90
8	5-Br	Me	H	—	3h/5h	4.5	97
9	—	—	—	H	4a/6a	2	89
10	—	—	—	4-OMe	4b/6b	3	95
11	—	—	—	4-Cl	4c/6c	3	81
12	—	—	—	3-Cl	4d/6d	2	96
13	—	—	—	2-Cl	4e/6e	2	92

^a Isolated yields after column chromatography.

cycloaddition step towards tetrazoles was optimized with regard to reaction temperature, concentration of the reactants, solvent composition and amounts of NaN_3 and ZnBr_2 promoter, to efficiently deliver the products in short times. The use of a hydroalcoholic medium turns the procedure eco-friendly and amenable to use in synthetic organic chemistry.

Acknowledgments

The authors are thankful to MCT/CNPq and CAPES, for financial support. T.S.K. also thanks CONICET.

Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.tetlet.2014.01.101>.

References and notes

- (a) Sadlej-Sosnowska, N. *J. Org. Chem.* **2001**, *66*, 8737–8743; (b) Mohite, P. B.; Bhaskar, V. H. *Int. J. Pharm. Tech Res.* **2011**, *3*, 1557–1566.
- (a) Ostrovskii, V. A.; Trifonov, R. E.; Popova, E. A. *Russ. Chem. Bull.* **2012**, *61*, 768–780; (b) Myznikov, L. V.; Hrabalek, A.; Koldobskii, G. I. *Chem. Heterocycl Comp.* **2007**, *43*, 1–9; (c) Herr, J. R. *Bioorg. Med. Chem.* **2002**, *10*, 3379–3393; (d) Sureshbabu, V. V.; Venkataramanarao, R.; Naik, S. A.; Chennakrishnareddy, G. *Tetrahedron Lett.* **2007**, *48*, 7038–7041.
- Gutmann, B.; Roduit, J.-P.; Roberge, D.; Kappe, C. O. *Angew. Chem., Int. Ed.* **2010**, *49*, 7101–7105.
- (a) Nasrallahzadeh, M.; Bayat, Y.; Habibi, D.; Moshaei, S. *Tetrahedron Lett.* **2009**, *50*, 4435–4438; (b) Sureshbabu, V. V.; Vasantha, B.; Hemantha, H. P. *Synthesis* **2011**, 1447–1455; (c) Heravi, M. M.; Fazeli, A.; Oskooie, H. A.; Beheshtiha, Y. S.; Valizadeh, H. *Synlett* **2012**, 2927–2930; (d) Roh, J.; Vávrová, K.; Hrabálek *Eur. J. Org. Chem.* **2012**, 6101–6118; (e) Venkateshwari, G.; Premalatha, A.; Rajanna, K. C.; Saiprakash, P. K. *Synth. Commun.* **2009**, *39*, 4479–4485.
- (a) Hajra, S.; Sinha, D.; Bhowmick, M. *J. Org. Chem.* **2007**, *72*, 1852–1855; (b) Laha, J. K.; Cuny, G. D. *Synthesis* **2008**, 4002–4006; (c) Curran, D. P.; Hadida, S.; Kim, S.-Y. *Tetrahedron* **1999**, *55*, 8997–9006; (d) Billot, X.; Chateauneuf, A.; Chauret, N.; Denis, D.; Greig, G.; Mathieu, M.-C.; Metters, K. M.; Slipetz, D. M.; Young, R. N. *Bioorg. Med. Chem. Lett.* **2003**, *13*, 1129–1132; (e) Aureggi, V.; Sedelmeier, G. *Angew. Chem., Int. Ed.* **2007**, *46*, 8440–8444; (f) Sedelmeier, G. US Patent 7,943,647, 2011; (g) Keith, J. M. *J. Org. Chem.* **2006**, *71*, 9540–9543.
- (a) Das, B.; Reddy, C. R.; Kumar, D. N.; Krishnaiah, M.; Narendar, R. *Synlett* **2010**, 391–394; (b) Cantillo, D.; Gutmann, B.; Kappe, C. O. *J. Am. Chem. Soc.* **2011**, *133*, 4465–4475; (c) Roh, J.; Artamonova, T. V.; Vávrová, K.; Koldobskii, G. I.; Hrabálek, A. *Synthesis* **2009**, 2175–2178.
- (a) Bazile, Y.; Bras, J. P.; De Cointet, P.; Nanthavong, S.; Pigerol, C.; Broll, M.; Eymard, P.; Werbenec, J. P.; Fournier, J. *Eur. J. Med. Chem.* **1977**, *12*, 525–530; (b) Finnegan, W. G.; Henry, R. A.; Lofquist, R. *J. Am. Chem. Soc.* **1958**, *80*, 3908–3911; (c) Jin, T.; Kitahara, F.; Kamijo, S.; Yamamoto, Y. *Tetrahedron Lett.* **2008**, *49*, 2824–2827; (d) Matthews, D. P.; Green, J. E.; Shuker, A. *J. J. Comb. Chem.* **2000**, *2*, 19–23.
- Demko, Z. P.; Sharpless, K. B. *Org. Lett.* **2001**, *3*, 4091–4094.
- (a) Demko, P. Z.; Sharpless, K. B. *J. Org. Chem.* **2001**, *66*, 7945–7950; (b) Himo, F.; Demko, Z. P.; Noodleman, L.; Sharpless, K. B. *J. Am. Chem. Soc.* **2002**, *124*, 12210–12216; (c) Demko, Z. P.; Sharpless, K. B. *Org. Lett.* **2002**, *4*, 2525–2527.
- (a) Lang, L.; Li, B.; Liu, W.; Jiang, L.; Xu, Z.; Yin, G. *Chem. Commun.* **2010**, 448–450; (b) Lakshmi Kantam, M.; Shiva Kumar, K. B.; Phani Raja, K. J. *Mol. Catal. A.* **2006**, *247*, 186–188; (c) Schulz, M. J.; Coatz, S. J.; Hlasta, D. J. *Org. Lett.* **2004**, *6*, 3265–3268; (d) Rostamizadeh, S.; Ghaieni, H.; Aryan, R.; Amani, A. *Chin. Chem. Lett.* **2009**, *20*, 1311–1314; (e) Kanakaraju, S.; Prasanna, B.; Chandramouli, G. V. P. *J. Chem.* **2013**, art id 104690; (f) Nandre, K. P.; Salunke, J. K.; Nandre, J. P.; Patil, V. S.; Borse, A. U.; Bhosale, S. V. *Chin. Chem. Lett.* **2012**, *23*, 161–164.
- (a) Newman, A. A. *Chemistry and Biochemistry of Thiocyanic Acid and its Derivatives*, 1st ed.; Academic Press: USA, 1975; (b) Khazei, A.; Zolfogol, M. A.; Safaiee, M.; Mohammad, M.; Donyadari, E.; Shiri, S. M.; Kruger, H. G. *Catal. Commun.* **2012**, *26*, 34–38.
- (a) Chakrabarty, M.; Sarkar, S. *Tetrahedron Lett.* **2003**, *44*, 8131–8133; (b) Murthy, Y. L. N.; Govindh, B.; Diwakar, B. S.; Nagalakshmi, K.; Venu, R. *Iran. Chem. Soc.* **2011**, *8*, 292–297; (c) Das, B.; Kumar, A. S. *Synth. Commun.* **2010**, *40*, 337–341; (d) Raju, R. M.; Lenin, R. *Lett. Org. Chem.* **2010**, *7*, 392–395.
- (a) Mahajan, U. S.; Akamanchi, K. G. *Synth. Commun.* **2009**, *39*, 2674–2682; (b) Wu, J.; Wu, G.; Wu, L. *Synth. Commun.* **2008**, *38*, 2367–2373; (c) Nair, V.; George, T. G.; Nair, L. G.; Panicker, S. B. *Tetrahedron Lett.* **1999**, *40*, 1195–1196; (d) Yadav, J. S.; Reddy, B. V. S.; Shubashree, S.; Sadashiv, K. *Tetrahedron Lett.* **2004**, *45*, 2951–2954; (e) Pan, X. Q.; Lei, M. Y.; Zou, J. P.; Zhang, W. *Tetrahedron Lett.* **2009**, *50*, 347–349; (f) Yadav, J. S.; Reddy, B. V. S.; Krishna, A. D.; Reddy, C. S.; Narsaiah, A. V. *Synthesis* **2005**, 961–964; (g) Memarian, H. R.; Mohammadpoor-Baltork, I.; Nikoofar, K. *Ultrason. Sonochem.* **2008**, *15*, 456–462; (h) Khazei, A.; Zolfogol, M. A.; Mokhlesi, M.; Panah, F. D.; Sajjadifar, S. *Helv. Chim. Acta* **2012**, *95*, 106–114; (i) Wu, G.; Liu, Q.; Shen, Y.; Wu, W.; Wu, L. *Tetrahedron Lett.* **2005**, *46*, 5831–5834.
- (a) da Silva, F. M. C.; dos Santos, J. C.; Campos, J. L. O.; Mafud, A. C.; Polikarpov, I.; Figueira, A. C. M.; Nascimento, A. S. *Bioorg. Med. Chem. Lett.* **2013**, *23*, 5795–5802; (b) Kaplancikli, Z. A.; Altintop, M. D.; Turan-Zitouni, G.; Ozdemir, A.; Can, O. D. *J. Enzyme Inhibit. Med. Chem.* **2012**, *27*, 275–280; (c) Altintop, M. D.; Ozdemir, A.; Turan-Zitouni, G.; Ilgin, S.; Atli, Ö.; Iscan, G.; Kaplancikli, Z. A. *Eur. J. Med. Chem.* **2012**, *58*, 299–307; (d) Adamcic, J.; Waisser, K.; Kunes, J.; Kauštová, J. *Arch. Pharm. Chem. Life Sci.* **2005**, *338*, 385–389.
- (a) Nichick, M. N.; Voitekhovich, S. V.; Shavel, A.; Lesnikovich, A. I.; Ivashkevich, O. A. *Polyhedron* **2009**, *28*, 3138–3142; (b) Wincott, F.; DiRenzo, A.; Shaffer, C.; Grimm, S.; Tracz, D.; Workman, C.; Sweedler, D.; González, C.; Scarine, S.; Usman, N. *Nucleic Acids Res.* **1995**, *23*, 2677–2684; (c) Tsou, D.; Harnpel, A.; Andrus, A.; Vinayak, R. *Nucleosides Nucleotides* **1995**, *14*, 1481–1492.
- (a) Lieber, E.; Enkoji, T. *J. Org. Chem.* **1961**, *26*, 4472–4479; (b) Patonay, T.; Léval, A. *Arch. Pharm. (Weinheim)* **1994**, *327*, 181–186.
- (a) Mueller, S.; Welz, R. *Tetrahedron Lett.* **2002**, *43*, 795–797; (b) LeBlanc, B. W.; Jursic, B. S. *Synth. Commun.* **1998**, *28*, 3591–3599.
- (a) Disli, A.; Salman, M. *Russ. J. Org. Chem.* **2009**, *45*, 151–153; (b) Kolo, K.; Sajadi, S. M. *Lett. Org. Chem.* **2013**, *10*, 688–692.
- (a) Prieto, M.; Zurita, E.; Rosa, E.; Muñoz, L.; Lloyd-Williams, P.; Giralt, E. *J. Org. Chem.* **2004**, *69*, 6812–6820; (b) Roxburgh, C. J.; Sammes, P. G.; Abdullah, A. *Dyes Pigments* **2009**, *82*, 226–237; (c) Samsoniya, S. A.; Barbakadze, N. N.; Chikvaidze, I. Sh. *ARKIVOC* **2012**, *143*–154; (d) Roy, S.; Eastman, A.; Gribble, G. W. *Tetrahedron* **2006**, *62*, 7838–7845; (e) Yang, X.; Xing, H.; Zhang, Y.; Lai, Y.; Zhang, Y.; Jiang, Y.; Ma, D. *Chin. J. Chem.* **2012**, *30*, 875–880.
- Silveira, C. C.; Fortes, M. P.; Mendes, S. R. *Curr. Org. Chem.* **2012**, *16*, 1540–1548.
- Typical procedure for the synthesis of the thiocyanates: A magnetically stirred mixture of the indole (1.0 mmol) and NH_4SCN (114 mg, 1.5 mmol) in MeOH (10 mL), was treated with Oxone (921 mg, 1.5 mmol), and the reaction was stirred at room temperature until complete consumption of the starting material (TLC). Water (30 mL) was added and the products were extracted with EtOAc (3 × 10 mL). The organic phase was dried (MgSO_4), concentrated under reduced pressure and the residue was purified by chromatography (hexane/EtOAc, 90:10).
- Alerman, M.; Hallberg, A. *J. Org. Chem.* **2000**, *65*, 7984–7989.
- Typical procedure for the synthesis of the 5-sulfonyl tetrazoles: A stirred mixture of the thiocyanate (1 mmol, 1.0 equiv) Na_3 (1.2 equiv) and ZnBr_2 (1.0 equiv) in 2-PrOH/H₂O (1:1, 6 mL) was refluxed until complete consumption of the starting material (2–4.5 h). After cooling to room temperature, the reaction was acidified with 1 M HCl (10 mL) and diluted with EtOAc (15 mL). The aqueous phase was extracted with EtOAc (2 × 20 mL) and the combined organic layers were dried (MgSO_4) and concentrated under reduced pressure. The residue was purified by column chromatography (hexane/EtOAc to EtOAc/MeOH 90:10).
- Characterization of selected compounds: ¹H and ¹³C NMR were run at 400 and 100 MHz, respectively, with the samples dissolved in DMSO-d₆. 3-((1H-Tetrazol-5-yl)thio)-1H-indole (**5a**). Beige solid, mp: 210.4–211.6 °C. IR (KBr, v): 3269, 2336, 1653, 1508, 1310, 1042, 746 and 604 cm⁻¹. ¹H NMR δ: 11.83 (s, 1H), 7.91 (d, 1H, *J* = 2.3 Hz), 7.55 (d, 1H, *J* = 8.0 Hz), 7.49 (d, 1H, *J* = 7.8 Hz), 7.23–7.20 (m, 1H) and 7.14–7.10 (m, 1H). ¹³C NMR δ: 155.3, 136.6, 133.1, 128.3, 122.3, 120.4, 117.9, 112.9 and 94.0. EI-MS (*m/z*, rel int, %): 218 [(M+1)⁺, 2], 217 (M⁺, 150), 160 (29), 148 (100), 89 (19) and 77 (19). Anal. Calcd: C, 49.76; H, 3.25. Found: C, 50.09; H, 3.55. 3-((1H-Tetrazol-5-yl)thio)-5-bromo-1H-indole (**5b**). Beige solid, mp: 209.5–211.3 °C. IR (KBr, v): 3274, 2364, 1566, 1309, 1217, 1109, 1033, 795 and 746 cm⁻¹. ¹H NMR δ: 12.00 (s, 1H), 7.93 (d, 1H, *J* = 2.5 Hz), 7.64 (s, 1H), 7.49 (d, 1H, *J* = 6.6 Hz) and 7.31 (dd, 1H, *J* = 8.6 Hz, 0.9 Hz). ¹³C NMR δ: 155.1, 135.4, 134.6, 130.4, 125.0, 120.3, 114.5, 113.3 and 94.2. EI-MS (*m/z*, rel int, %): 296 [(M+2)⁺, 34], 295 [(M+1)⁺, 4], 294 (M⁺, 31), 239 (64), 227 (97), 225 (100), 173 (40), 148 (37), 120 (52), 114 (16) and 45 (34). HRMS obsd *m/z*: 317.9429; C₉H₆BrN₅SNa [M+Na]⁺ requires *m/z* 317.9425. 3-((1H-Tetrazol-5-yl)thio)-5-(*p*-tolyl)-1H-indole (**5c**). Brown solid, mp: 213.3–215.5 °C. IR (KBr, v): 3263, 2331, 1502, 1473, 1308, 1108, 1036 and 799 cm⁻¹. ¹H NMR δ: 11.81 (s, 1H), 7.90 (s, 1H), 7.65 (s, 1H), 7.58 (*d*, *J* = 8.4 Hz, 1H), 7.52–7.50 (m, 3H), 7.23 (d, 2H, *J* = 7.4 Hz), 2.33 (s, 3H). ¹³C NMR δ: 154.9, 137.9, 135.6, 135.3, 133.2, 132.7, 128.9, 128.5, 126.1, 121.2, 115.1, 112.4, 94.3 and 20.1. EI-MS (*m/z*, rel int, %): 308 [(M+1)⁺, 2], 307 (M⁺, 11), 277 (18), 238 (44), 173 (100), 11 (21), 77 (9) and 45 (21). Anal. Calcd: C, 62.58; H, 4.26. Found: C, 62.68; H, 4.68. 3-((1H-Tetrazol-5-yl)thio)-2-methyl-1H-indole (**5d**). Beige solid, mp: 184.5–185.7 °C. IR (KBr, v): 3339, 3052, 2332, 1546, 1518, 1314, 1035 and 742 cm⁻¹. ¹H NMR δ: 11.68 (s, 1H), 7.43–7.39 (m, 2H), 7.16–7.12 (m, 1H), 7.09–7.06 (m, 1H) and 2.52 (s, 3H). ¹³C NMR δ: 155.2, 143.1, 135.5, 129.2, 121.4, 120.0, 117.1, 111.3, 91.3 and 11.6. EI-MS (*m/z*, rel int, %): 233 [(M+2)⁺, 2], 233 [(M+1)⁺, 5], 231 (M⁺, 34), 174 (33), 162 (100), 130 (32), 118 (67) and 77 (27). HRMS obsd *m/z*: 254.0471; C₁₀H₉N₅SNa [M+Na]⁺ requires *m/z* 254.0476. 3-((1H-Tetrazol-5-yl)thio)-1-methyl-1H-indole (**5e**). Beige solid, mp: 179.9–181.4 °C. IR (KBr, v): 3394, 3110, 2331, 1517, 1243, 1028 and 744 cm⁻¹. ¹H NMR δ: 7.87 (s, 1H), 7.55 (d, *J* = 8.2 Hz, 1H), 7.49 (d, *J* = 7.8 Hz, 1H), 7.29–7.26 (m, 1H), 7.18–7.14 (m, 1H) and 3.85 (s, 3H). ¹³C NMR δ: 155.2, 137.1, 136.6, 128.6, 122.2, 120.4, 118.0, 110.5, 92.9 and 32.7. EI-MS (*m/z*, rel int, %): 231 (M⁺, 4), 174 (27), 162 (100), 118 (15) and 77 (16). Anal. Calcd: C, 51.93; H, 3.92. Found: C, 51.99; H, 3.99. 3-((1H-Tetrazol-5-yl)thio)-1-octyl-1H-indole (**5f**). Yellow solid, mp: 95.4–96.9 °C. IR (KBr, v): 3444, 2953, 2923, 2852, 2713, 1613, 1508, 1492, 1457, 1299, 1165, 1020 and 738 cm⁻¹. ¹H NMR δ: 7.91 (s, 1H), 7.59

(d, 1H, $J = 8.1$ Hz), 7.47 (d, 1H, $J = 7.7$ Hz), 7.27–7.23 (m, 1H), 7.16–7.12 (m, 1H), 4.22 (t, 2H, $J = 6.8$ Hz), 1.85–1.75 (m, 2H), 1.28–1.22 (m, 10H) and 0.85–0.83 (m, 3H). ^{13}C NMR δ : 155.1, 136.4, 135.5, 128.6, 122.1, 120.3, 118.1, 110.6, 93.0, 45.8, 30.9, 29.3, 28.3, 25.9, 21.7 and 13.6. EI-MS (m/z , rel int.-%): 331 [(M+2) $^+$, 2], 330 [(M+1) $^+$, 7], 329 (M $^+$, 33), 272 (57), 260 (59), 162 (100), 130 (28), 77 (8), 57 (26) and 41 (50). Anal. Calcd: C, 61.97; H, 7.04. Found: C, 61.74; H, 7.23. 3-((1*H*-Tetrazol-5-yl)thio)-1-phenyl-1*H*-indole (**5g**). Yellow solid, mp: 153.1–154.3 °C. IR (KBr, ν): 3033, 2331, 1598, 1499, 1457, 1231, 1017, 746 and 695 cm^{-1} . ^1H NMR δ : 8.22 (d, $J = 0.9$ Hz, 1H), 7.66–7.56 (m, 6H), 7.49–7.45 (m, 1H), 7.33–7.29 (m, 1H) and 7.26–7.22 (m, 1H). ^{13}C NMR δ : 154.9, 137.9, 136.0, 135.5, 129.8, 129.1, 127.3, 124.2, 123.5, 121.6, 118.6, 111.1 and 96.9. EI-MS (m/z , rel int.-%): 294 [(M+1) $^+$, 5], 293 (M $^+$, 29), 236 (36), 224 (100), 121 (21) and 77 (40). HRMS obsd m/z : 316.0614; $\text{C}_{15}\text{H}_{11}\text{N}_5\text{SNa}$ [M+Na] $^+$ requires m/z 316.0633. 3-((1*H*-Tetrazol-5-yl)thio)-5-bromo-1-methyl-1*H*-indole (**5h**). Beige solid, mp: 205.5–207.0 °C. IR (KBr, ν): 3394, 1509, 1465, 1242, 795 and 614 cm^{-1} . ^1H NMR δ : 7.87 (s, 1H), 7.62 (d, $J = 1.7$ Hz, 1H), 7.52 (d, $J = 8.7$ Hz, 1H), 7.36 (dd, $J = 8.7$ Hz, 1.8 Hz, 1H) and 3.82 (s, 3H). ^{13}C NMR δ : 155.2, 137.6, 135.8, 130.6, 124.7, 120.5, 113.2, 112.8, 94.3 and 33.0. EI-MS (m/z , rel int.-%): 310 [(M+2) $^+$, 35], 309 [(M+1) $^+$, 5], 308 (M $^+$, 35), 253 (48), 241 (97), 239 (100), 162 (24), 117 (32) and 77 (7). HRMS obsd m/z : 331.9575; $\text{C}_{10}\text{H}_8\text{BrN}_5\text{SNa}$ [M+Na] $^+$ requires m/z 331.9581. 5-((1-Phenyl-1*H*-pyrrol-2-yl)thio)-1*H*-tetrazole (**6a**). Yellow solid, mp: 160.1–161.4 °C. IR (KBr, ν): 3048, 2366,

- 1595, 1498, 1322, 1038, 766 and 740 cm^{-1} . ^1H NMR δ : 7.45–7.34 (m, 6H), 6.82 (dd, $J = 3.6$ Hz, 1.6 Hz, 1H) and 6.40–6.38 (m, 1H). ^{13}C NMR δ : 155.0, 138.3, 128.7, 128.1, 127.7, 126.0, 121.6, 111.4 and 109.9. EI-MS (m/z , rel int.-%): 244 [(M+1) $^+$, 2], 243 (M $^+$, 6), 242 (37), 185 (34), 174 (45), 173 (100), 172 (48), 130 (16), 115 (19), 77 (65), 70 (26) and 51 (39). HRMS obsd m/z : 266.0465; $\text{C}_{11}\text{H}_9\text{N}_5\text{SNa}$ [M+Na] $^+$ requires m/z 266.0476. 5-((1-(4-Methoxyphenyl)-1*H*-pyrrol-2-yl)thio)-1*H*-tetrazole (**6b**). White solid, mp: 210.0–211.4 °C. IR (KBr, ν): 3012, 2360, 1517, 1250, 1039, 837 and 740 cm^{-1} . ^1H NMR δ : 7.27–7.24 (m, 3H), 6.96 (d, $J = 8.9$ Hz, 2H), 6.78 (dd, $J = 3.6$ Hz, 1.7 Hz, 1H), 6.39–6.35 (m, 1H) and 3.76 (s, 3H). ^{13}C NMR δ : 158.6, 155.1, 131.3, 128.4, 127.5, 121.2, 113.9, 111.5, 109.6 and 55.2. EI-MS (m/z , rel int.-%): 275 [(M+2) $^+$, 2], 274 [(M+1) $^+$, 6], 273 (M $^+$, 37), 216 (20), 204 (100), 173 (37), 92 (15) and 77 (20). Anal. Calcd: C, 52.73; H, 4.06. Found: C, 52.86; H, 4.31. 5-((1-(4-Chlorophenyl)-1*H*-pyrrol-2-yl)thio)-1*H*-tetrazole (**6c**). Orange solid, mp: 171.2–173.1 °C. IR (KBr, ν): 3129, 3010, 2362, 1494, 1311, 1094, 837 and 739 cm^{-1} . ^1H NMR δ : 7.50 (d, $J = 8.8$ Hz, 2H), 7.40 (d, $J = 8.8$, 2H), 7.35 (dd, 1H, $J = 3.0$ Hz, 1.8 Hz), 6.82 (dd, $J = 3.7$, 1.8, 1H) and 6.40 (dd, 1H, $J = 3.7$ Hz, 3.0 Hz). ^{13}C NMR δ : 155.0, 137.2, 132.4, 128.8, 128.3, 127.9, 122.0, 111.6 and 110.2. EI-MS (m/z , rel int.-%): 279 [(M+2) $^+$, 6], 278 [(M+1) $^+$, 2], 277 (M $^+$, 18), 219 (19), 173 (100), 111 (13) and 75 (19). HRMS obsd m/z : 300.0074; $\text{C}_{11}\text{H}_8\text{ClN}_5\text{SNa}$ [M+Na] $^+$ requires m/z 300.0087.
25. (a) Himo, F.; Demko, Z. P.; Noddleman, L.; Sharpless, B. K. *J. Am. Chem. Soc.* **2003**, 125, 9983–9987.